From the session 2023-24, the textbooks are rationalised under the new National Education Policy 2020. This **Sanjiv Refresher** is completely based on the new rationalised textbook.

Sanjiv Refresher Mathematics

Based on the Latest CBSE Syllabus and NCERT Text Book

Class - IX

Main Features

- Strictly designed as per the CBSE syllabus 2023-24 and latest NCERT text book edition.
- Detailed Explanation of Important Concepts for better understanding.
- Solved Additional Examples.
- NCERT Textbook Exercises with Solutions.
- Exercises for Extra Practice.
- Objective Type Questions, Very Short Answer Type Questions and Case Study Based Questions as per the latest CBSE syllabus 2023-24.
- Important Questions from NCERT Exemplar.

Publisher :



Jaipur



Price : ₹ 300/-

Published by : SANJIV PRAKASHAN

Dhamani Market, Chaura Rasta, **JAIPUR - 302003** email : sanjeevprakashanjaipur@gmail.com website : www.sanjivprakashan.com

© Publisher

Laser Type Setting : Sanjiv Prakashan (DTP Department) Jaipur

DISCLAIMER

Every effort has been made to remove the mistakes and omissions in this book. In case of any suggestion or any error found, send an email or post a letter to us at the following address :

Email : sanjeevprakashanjaipur@gmail.com

Postal Address : Publication Department

Sanjiv Prakashan,

Dhamani Market, Chaura Rasta,

Jaipur-302003

Your suggestions shall be taken care of in our next edition.

Though all the precautions have been taken in publishing this book yet for any mistake the author, publisher or printer is not liable. All disputes are subjected to Jaipur Jurisdiction only.

© Rights are reserved. No part of this publication may be produced in any form, by photocopy, microfilm, xerography, recording or otherwise without the prior written permission of the publisher. Any breach will entail legal action and prosecution without further notice.

Contents ||

1.	Number Systems	1 - 40
2.	Polynomials	41 - 75
3.	Coordinate Geometry	76 - 89
4.	Linear Equations in Two Variables	90 - 103
5.	Introduction to Euclid's Geometry	104 - 119
6.	Lines and Angles	120 - 139
7.	Triangles	140 - 176
8.	Quadrilaterals	177 - 208
9.	Circles	209 - 250
10.	Heron's Formula	251 - 263
11.	Surface Areas and Volumes	264 - 289
12.	Statistics	290 - 325

(iii)

NUMBER SYSTEMS

1.1. Introduction

examp

- **1.2. Irrational Numbers**
- 1.3. Real Numbers and their Decimal Expansions

08

- 1.4. Operations on Real Numbers
- 1.5. Laws of Exponents for Real Numbers

1.1. Introduction

- In earlier classes, we have learnt about number system, number line and how to represent various types of numbers on it.
- A number is a mathematical object used to count and measure, etc.
- Natural numbers are used for counting, *i.e.* 1, 2, 3, 4,..... the collection of natural numbers is denoted by N.

 $N = \{1, 2, 3, 4, 5, \dots\}$

• Whole numbers are all natural numbers together with zero. The collection of whole numbers is denoted by *W*.

 $W = \{0, 1, 2, 3, 4, \dots\}$

• Integers are all whole numbers and negative of natural numbers. The collection of integers is denoted by *Z* (Zahlen) or *I*.

 $Z \text{ or } I = \{\dots, -4, -3, -2, -1, 0, 1, 2, 3, 4, \dots\}$

• **Rational numbers :** Any number that can be expressed in the form $\frac{p}{q}$, where p and q are both integers and $q \neq 0$ is called a rational number.

The word 'rational' comes from the word 'ratio'. Thus every rational number can be written as a ratio of two integers. Set of rational numbers can be expressed as Q such that :

$$Q = \left\{ \frac{p}{q} : p, q \text{ are integers and } q \neq 0 \right\}$$

Note that every integer (positive, negative or zero) can be written in $\frac{p}{q}$ form where q = 1.

For example :

$$5 = \frac{5}{1}, -7 = \frac{-7}{1}, 0 = \frac{0}{1}$$

Sanjiv Refresher Mathematics—IX

Hence, every integer is a rational number. We know that the rational numbers do not have unique representation in the form $\frac{p}{q}$, where p and q are both integers

and $q \neq 0$. For example : $\frac{1}{2} = \frac{2}{4} = \frac{3}{6} = \frac{7}{14}$, etc. In fact these are equivalent rational numbers. However, if we write a rational number in the form $\frac{p}{q}$, where p and q are both integers and $q \neq 0$ and p and q have no common factors except 1, then among the infinitely many rational numbers equivalent to $\frac{1}{2}$, we choose $\frac{1}{2}$ to represent all of them on the number line.

Thus, a rational number can be uniquely expressed as $\frac{p}{q}$ where p and q are both integers, $q \neq 0$ and p and q have no common factors except 1, *i.e.* p and q are co-prime. It is called in the lowest terms.

Representation of rational numbers : We know that natural numbers, whole numbers can be represented on a number line. We can also represent rational numbers on the same number line.

Here the length OA represents unit length. As we can see in the figure above, every integer has been represented by one and only one point on the number line l.

Next, we consider the representation of rational numbers on the line 'l'. Take one-half of the unit length and mark points on 'l' on both sides of *O*; these points will represent the numbers $\frac{1}{2}, \frac{2}{2}, \frac{3}{2}$ and $\frac{-1}{2}, \frac{-2}{2}, \frac{-3}{2}$.

Similarly, take one third of the unit length and mark points on \mathcal{V} in both sides of O; these points will represent the numbers $\frac{1}{3}, \frac{2}{3}, \frac{3}{3}, \frac{4}{3}, \dots$ and $\frac{-1}{3}, \frac{-2}{3}, \frac{-3}{3}, \dots$ and so on.

- All natural numbers, whole numbers and integers are rational numbers.
- There are infinitely many rational numbers between any two given rational numbers.

Thus every rational number has been represented by one and only one point on the line l'.

- The collection of rational numbers is ordered, *i.e.* if a, b are any two rational numbers, then either a < b or a > b or a = b.
- Also, if a < b, then *n* rational numbers lying between 'a' and 'b' are (a + d), (a + 2d), (a + 3d), (a + nd) where, $d = \frac{b-a}{n+1}$

2

Number Systems

For example : If we have to find 4 rational numbers between -1 and -2 we have a < b number are (a + d); (a + 2d); (a + 3d); (a + 4d).

Here a = -2, b = -1

$$d = \frac{b-a}{n+1} = \frac{-1-1(-2)}{4+1} = \frac{-1+2}{5} = \frac{1}{5}$$

Rational numbers are

Where,

$$\left(-2+\frac{1}{5}\right);\ \left(-2+\frac{2}{5}\right);\ \left(-2+\frac{3}{5}\right);\ \left(-2+\frac{4}{5}\right) \Rightarrow \frac{-9}{5},\ \frac{-8}{5},\ \frac{-7}{5},\ \frac{-6}{5}$$

Hence 4 rational numbers between -1 and -2 are (-1), $\frac{-9}{5}$, $\frac{-8}{5}$, $\frac{-7}{5}$, $\frac{-6}{5}$, (-2)

Another method of representing rational numbers between any two rational numbers :

Let *a* and *b* be any two rational numbers, then to find a rational number between *a* and *b*. Add *a* and *b* and divide the sum by 2, *i.e.* if *a*, *b* are any two different rational numbers, then $\frac{a+b}{2}$ is a rational number and it lies between them, *i.e.* $a < \frac{a+b}{2} < b$. Continuing this process, we find that there are infinitely many rational numbers between two rational numbers.

For example :
$$\frac{\frac{1}{3} + \frac{2}{7}}{2} = \frac{13}{42}$$
 is a rational number which lies between $\frac{1}{3}$ and $\frac{2}{7}$.

One more method of finding 'n' rational numbers between any two rational numbers :

To insert '*n*' rational numbers between two rational numbers we proceed as under. **Case I**: When the given rational numbers have the same denominators, multiply the numerator and denominator of each rational number by (n + 1).

Case II : When the given rational numbers have different denominators.

(i) Find the LCM of denominators.

(ii) Change the given rational numbers to equivalent rational numbers with the LCM as their common denominators.

(iii) If necessary, multiply the numerator and denominator of each rational number by (n + 1).

23

 $, \frac{1}{28}$

For example :

Case I: Insert 3 rational numbers between $\frac{5}{7}$ and $\frac{6}{7}$.

Multiply and divide both by (n + 1), *i.e.* 3 + 1 = 4

$$\frac{5}{7} \times \frac{4}{4} = \frac{20}{28}$$

$$\frac{6}{7} \times \frac{4}{4} = \frac{24}{28}$$
Now 3 rational numbers between $\frac{20}{28}$ and $\frac{24}{28}$ are $\frac{21}{28}$, $\frac{22}{28}$

3

Sanjiv Refresher Mathematics—IX

Case II : Insert 3 rational numbers between $\frac{2}{9}$ and $\frac{3}{7}$. LCM of 7 and 9 = 63.

$$\frac{3}{7} \times \frac{9}{9} = \frac{27}{63}$$
$$\frac{2}{9} \times \frac{7}{7} = \frac{14}{63}$$
$$\therefore 3 \text{ rational numbers between } \frac{14}{63} \text{ and } \frac{27}{63} \text{ are } \frac{15}{63}, \frac{16}{63}, \frac{17}{63}.$$



Example 1. Find nine rational numbers between 0 and 0.1. Solution : Here, 0.1 > 0. So, let a = 0, b = 0.1 and n = 9

Now,
$$d = \frac{b-a}{n+1} = \frac{0.1-0}{9+1} = \frac{0.1}{10} = 0.01$$

So, the nine rational numbers between 0 and 0.1 are : (x + d), (x + 2d), (x + 3d), (x + 4d), (x + 5d), (x + 6d), (x + 7d), (x + 8d) and (x + 9d), *i.e.* (0 + 0.01), (0 + 2 × 0.01), (0 + 3 × 0.01), (0 + 4 × 0.01), (0 + 5 × 0.01), (0 + 6 × 0.01), (0 + 7 × 0.01), (0 + 8 × 0.01) and (0 + 9 × 0.01) = 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08 and 0.09 = $\frac{1}{100}, \frac{2}{100}, \frac{3}{100}, \frac{4}{100}, \frac{5}{100}, \frac{6}{100}, \frac{7}{100}, \frac{8}{100}$ and $\frac{9}{100}$ = $\frac{1}{100}, \frac{1}{50}, \frac{3}{100}, \frac{1}{25}, \frac{1}{20}, \frac{3}{50}, \frac{7}{100}, \frac{2}{25}$ and $\frac{9}{100}$ **NCERT Exercise 1.1**

1. Is zero a rational number? Can you write it in the form $\frac{p}{q}$, where p and q are integers and $q \neq 0$?

Solution : Yes, 0 is a rational number. 0 can be written in any of the following forms :

 $\frac{0}{1}, \frac{0}{-2}, \frac{0}{3}, \frac{0}{-4}$ and so on. Thus, 0 can be written as $\frac{p}{q}$, where p = 0 and q is any non-zero integer.

Hence, 0 is a rational number.

2. Find six rational numbers between 3 and 4.

Solution : We know that between two rational numbers *a* and *b*, such that a < b. There is always a rational number $\frac{a+b}{2}$, where $a < \frac{a+b}{2} < b$. A rational number between 3 and 4 is $\frac{1}{2}(3+4)$, *i.e.* $\frac{7}{2}$.

4